



1


The Knee Joint

[click here](#)

Main Menu

Copyright HandsOn Therapy Schools 2009

K.7



Knee Joint

Knee Joint

- largest joint in body
- very complex
- primarily a hinge joint

Enlarged femoral condyles articulate on enlarged tibial condyles

Medial and lateral tibial condyles (medial & lateral tibial plateaus) - receptacles for femoral condyles

Tibia – medial

- bears most of weight




Bones

Fibula - lateral

serves as the attachment for knee joint structures

does not articulate with femur or patella

not part of knee joint



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

[Back](#)

[Next](#)


Bones

Patella

sesamoid (floating) bone

imbedded in quadriceps & patellar tendon

serves similar to a pulley in improving angle of pull, resulting in greater mechanical advantage in knee extension



Bones

Key bony landmarks

Superior and inferior patellar poles


Tibial tuberosity

Gerdy's tubercle

Medial & lateral femoral condyles

Upper anterior medial tibial surface

Head of fibula



Anterior view with patella removed

Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

Bones

- Three vasti muscles of quadriceps originate on proximal femur and insert on patellar superior pole
 - insertion is ultimately on tibial tuberosity via patella tendon
- Iliotibial tract of tensor fasciae latae inserts on Gerdy’s tubercle
- Sartorius, gracilis, and semitendinosus insert just below the medial condyle on upper anteromedial tibial surface
- Semimembranosus inserts posteromedially on medial tibial condyle
- Biceps femoris inserts primarily on fibula head
- Popliteus originates on lateral aspect of lateral femoral condyle
- Tibial collateral ligament originates on medial aspect of upper medial femoral condyle and inserts on medial tibial surface
- Fibula collateral originates on lateral femoral condyle very close to popliteus origin and inserts on fibular head



Joints

Knee joint proper (tibiofemoral joint)

classified as a ginglymus joint ; Sometimes referred to as trochoginglymus joint internal and external rotation occur during flexion ; Some argue for condyloid classification

Patellofemoral joint

arthrodial classification

gliding nature of patella on femoral condyles

Ligaments provide static stability

Quadriceps and hamstrings contractions produce dynamic stability

Articular cartilage surfaces on femur and tibia

Menisci form cushions between bones

attached to tibia

deepen tibial fossa

enhance stability

Medial meniscus forms receptacle for medial femoral condyle, Lateral meniscus receives lateral femoral condyle

Thicker on outside border and taper down very thin to inside border

Can slip about slightly, but held in place by various small ligaments

Medial meniscus - larger and more open C appearance

Lateral meniscus - closed C configuration

Either or both menisci may be torn in several different areas from a variety of mechanisms, resulting in varying degrees of problems

Tears often occur due significant compression and shear forces during rotation while flexing or extending during quick directional changes in running

Anterior and posterior cruciate ligaments

cross within knee between tibia and femur

vital in respectively maintaining anterior and posterior stability, as well as rotatory stability

Anterior cruciate ligament (ACL) injuries


one of most common serious injuries to knee

mechanism often involves noncontact rotary forces associated with planting and cutting, hyperextension, or by violent quadriceps contraction which pulls tibia forward on femur

[Back](#)

[Next](#)

Joints



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.



Joints

Posterior cruciate ligament (PCL) injuries

- not often injured

- mechanism of direct contact with an opponent or playing surface

Fibular (lateral) collateral ligament (LCL)

- infrequently injured

Tibial (medial) collateral ligament (MCL)

- maintains medial stability by resisting valgus forces or preventing knee from being abducted

- injuries occur commonly, particularly in contact or collision sports

- mechanism of teammate or opponent may fall against lateral aspect of knee or leg causing medial opening of knee joint and stress to medial ligamentous structures

Synovial cavity

- supplies knee with synovial fluid

- lies under patella and between surfaces of tibia and femur

- "capsule of the knee"

Infrapatellar fat pad

- just posterior to patellar tendon

- an insertion point for synovial folds of tissue known as "plica"

- an anatomical variant that may be irritated or inflamed with injuries or overuse of the knee

Bursae

- more than 10 bursae in and around knee

- some are connected to synovial cavity

- they absorb shock or prevent friction


Joints

Extends to 180 degrees (0 degrees of flexion)

Hyperextension of 10 degrees or > not uncommon

Flexion occurs to about 140 degrees

With knee flexed 30 degrees or > ; internal rotation 30 degrees occurs ; external rotation 45 degrees occurs



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

Muscles

Knee “screws home” to fully extend due to the shape of medial femoral condyle

As knee approaches full extension tibia must externally rotate approximately 10 degrees to achieve proper alignment of tibial and femoral condyles


In full extension

close congruency of articular surfaces

no appreciable rotation of knee

During initial flexion from full extension

knee “unlocks” by tibia rotating internally, to a degree, from its externally rotated position to achieve flexion




Movements

Flexion

bending or decreasing angle between femur and leg, characterized by heel moving toward buttocks

Extension

straightening or increasing angle between femur and lower leg



Flexion

Extension

Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

Movements

External rotation

rotary movement of leg laterally away from midline


Internal rotation

rotary movement of lower leg medially toward midline

Neither will occur unless flexed 20-30 degrees or >



External rotation



Internal rotation

Muscles

Q angle

Central line of pull for entire quadriceps runs from ASIS to the center of patella

Line of pull of patella tendon runs from center of patella to center of tibial tuberosity


Angle formed by the intersection of these two lines at the patella is the Q angle

Normally, angle will be 15 degrees or less for males & 20 degrees or less in females

Generally, females have higher angles due to a wider pelvis

Higher Q angles generally predispose people in varying degrees to a variety of potential knee problems including lateral patellar subluxation or dislocation, patellar compression syndrome, chondromalacia, and ligamentous injuries

For people with above normal Q angles, it is particularly important to maintain high levels of strength & endurance in vastus medialis so as to counteract lateral pull of vastus lateralis




Muscles


Quadriceps muscle group

extends knee

located in anterior compartment of thigh

consists of 4 muscles: rectus femoris ; vastus lateralis ; vastus intermedius ; vastus medialis





Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

[Back](#)

[Next](#)

Muscles

Hamstring muscle group

responsible for knee flexion

located in posterior compartment of thigh

consists of 3 muscles ; semitendinosus - medial, internal rotator ; semimembranosus - medial, internal rotator ; biceps femoris - lateral, external rotator

Popliteus assist medial hamstrings in knee internal rotation

Two-joint muscles

most effective when either origin or insertion is stabilized to prevent movement in direction of the contacting muscle

To a degree, muscles are able to exert greater force when lengthened than when shortened

Hamstring muscles and rectus femoris are biarticular (two-joint) muscles : Ex. sartorius muscle

increases its total length and becomes a better flexor at knee when pelvis is rotated posteriorly and stabilized by abdominal muscles

exemplified by trying to flex knee and cross the legs in the sitting position

one usually leans backward to flex legs at knees

Football kicker invariably leans well backward to raise and fix the rectus femoris origin to make it more effective as a knee extensor

Gracilis, sartorius, and semitendinosus join together distally to form pes anserinus

attaches to anteromedial aspect of proximal tibia below the level of tibial tuberosity

Their attachment and posteromedially line of pull enable them to assist with knee flexion particularly once the knee is flexed and hip is externally rotated

Medial and lateral gastrocnemius heads attach posteriorly on medial and lateral femoral condyles

assist with knee flexion

Knee Joint Muscle Location


Anterior - primarily knee extension


Rectus femoris

Vastus medialis

Vastus intermedius

Vastus lateralis





Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

[Back](#)

[Next](#)

Knee Joint Muscle Location

Posterior - primarily knee flexion

Biceps femoris

Semimembranosus


Semitendinosus


Sartorius

Gracilis

Popliteus

Gastrocnemius





Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

[Back](#)

[Next](#)

Muscles

Quadriceps muscles - vital in jumping

- functions as a decelerator : when decreasing speed to change direction ; when coming down from a jump
- eccentric contraction during decelerating actions
- controls slowing of movements initiated in previous phases of the sports skill

Rectus femoris (two-joint), vastus medialis, vastus intermedius, vastus lateralis (largest)

- All attach to patella then to tibial tuberosity via patellar tendon
- All superficial and palpable except vastus intermedius (under rectus femoris)
- Strength or power may be indicated by vertical jump test
- Generally desired to be 25% to 33% stronger than hamstring group

Strength and endurance is essential for maintenance of patellofemoral stability


- often a problem
- quads are particularly prone to atrophy when injuries occur
- may be developed by resisted knee extension activities from a seated position
- functional weight bearing activities such as step-ups or squats are particularly useful for strengthening and endurance

Rectis Femoris Muscle

Flexion of hip

Extension of knee


Anterior pelvic rotation



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

Vastus Intermedius Muscle

Extension of knee




Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

[Back](#)

[Next](#)

Vastus Medialis Muscle

Extension of knee



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

[Back](#)


[Next](#)

Hamstring Muscle Group

Semitendinosus

Biceps femoris


Semimembranosus



Semitendinosus

(long head)
Biceps femois
(short head)

Semimembranosus



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

[Back](#)

[Next](#)

Hamstring Muscle Group

Semitendinosus

Biceps femoris

Semimembranosus

Hamstring muscle strains very common

“Running muscles” function in acceleration

Antagonists to quadriceps muscles at knee


Named for cordlike attachments at knee


All originate on ischial tuberosity of pelvis

Semitendinosus inserts on anteromedial tibia

Semimembranosus inserts on posteromedial tibia

Biceps femoris inserts on lateral tibial condyle and head of fibula





Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

Semitendinosus Muscle


Flexion of knee

Extension of hip

Internal rotation of hip

Internal rotation of flexed knee

Internal rotation of flexed knee



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

[Back](#)

[Next](#)

Semimembranosus Muscle


Flexion of knee

Extension of hip

Internal rotation of hip

Internal rotation of flexed knee

Posterior pelvis rotation



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

[Back](#)

[Next](#)

Biceps Femoris Muscle


Flexion of knee

Extension of hip

External rotation of hip

External rotation of flexed knee

Posterior pelvis rotation



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.


[Back](#)

[Next](#)

Popliteus Muscle

Flexion of knee

Internal rotation of flexed knee



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

Knee Extension


Agonist

Rectus Femoris

Vastus Lateralis

Vastus Intermedius

Vastus Medialis



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.


Knee Flexion

Agonist

Biceps Femoris (Long & Short Head)

Semitendinosus

Semimembranosus




Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

Knee External Rotation

Agonist

Biceps Femoris



Copyright 2007 McGraw-Hill Higher Education. All rights reserved.

To Test

Access Code: **BE3Y**

Please write down case sensitive code. You will be asked for it. This course is 6 CE.

Once you have successfully passed the test (70% correct), please email us at cehotschool@gmail.com. We will email you your CE certificate within 7 business days.